

Supplementary difference sets related to a certain class of complex spherical 2-codes

Makoto Araya*, Masaaki Harada† and Sho Suda‡

February 24, 2016

Dedicated to Professor Hiroshi Kimura on His 80th Birthday

Abstract

In this paper, we study skew-symmetric $2\text{-}\{v; r, k; \lambda\}$ supplementary difference sets related to a certain class of complex spherical 2-codes. A classification of such supplementary difference sets is complete for $v \leq 51$.

1 Introduction

Let $\Omega(d)$ denote the complex unit sphere in $\mathbb{C}^d$. For a finite set X in $\Omega(d)$, define

$$A(X) = \{x^*y \mid x, y \in X, x \neq y\},$$

where x^* is the transpose conjugate of a column vector x . A finite set X is called a *complex spherical 2-code* if $|A(X)| = 2$ and $A(X)$ contains an imaginary number. A complex spherical 2-code X with $A(X) = \{\alpha, \bar{\alpha}\}$ has the structure of a tournament (X, E) , where $E = \{(x, y) \in X \times X \mid x^*y = \alpha\}$ [12]. We say that the tournament (X, E) is attached to the complex spherical 2-code X .

*Department of Computer Science, Shizuoka University, Hamamatsu 432–8011, Japan. email: araya@inf.shizuoka.ac.jp

†Research Center for Pure and Applied Mathematics, Graduate School of Information Sciences, Tohoku University, Sendai 980–8579, Japan. email: mharada@m.tohoku.ac.jp.

‡Department of Mathematics Education, Aichi University of Education, Kariya 448–8542, Japan. email: suda@aecc.aichi-edu.ac.jp

Theorem 1 (Nozaki and Suda [12, Theorem 4.8]). *Let X be a complex spherical 2-code in $\Omega(d)$. Let A be the adjacency matrix of the tournament G attached to X .*

- (1) $|X| \leq 2d + 1$ if d is odd, and $|X| \leq 2d$ if d is even.
- (2) $|X| = 2d + 1$ for odd d if and only if G is a doubly regular tournament.
- (3) $|X| = 2d$ for even d if and only if $I + A - A^T$ is a skew-Hadamard matrix, where I is the identity matrix and A^T denotes the transposed matrix of A .
- (4) $|X| = 2d$ for odd d if and only if one of the following occurs:
 - (a) A is obtained as the adjacency matrix of the induced subgraph of some doubly regular tournament by deleting a certain vertex.
 - (b) There exists a permutation matrix P such that

$$P(I + A - A^T)(I + A - A^T)^T P^T = \begin{pmatrix} \alpha I + \beta J & O \\ O & \alpha I + \beta J \end{pmatrix},$$

for some integers α, β with $\alpha \geq 2, \beta \geq 1$, where J denotes the all-one matrix and O denotes the zero matrix of appropriate size.

Doubly regular tournaments have been widely studied (see e.g., [8, 11, 13–15]). Skew-Hadamard matrices are a class of Hadamard matrices, which has been widely studied (see e.g., [2, 5, 6, 11, 13, 15, 18]). These motivate our investigation of matrices M satisfying the following conditions:

$$M \text{ is a } 2d \times 2d \text{ } (1, -1)\text{-matrix with } d \text{ odd,} \quad (1)$$

$$M - I = -(M - I)^T, \text{ that is, } M \text{ is skew-symmetric,} \quad (2)$$

$$MM^T = \begin{pmatrix} \alpha I + \beta J & O \\ O & \alpha I + \beta J \end{pmatrix} \text{ for some integers } \alpha, \beta \text{ with } \alpha \geq 2, \beta \geq 1. \quad (3)$$

In this paper, with this motivation, we study skew-symmetry for 2- $\{v; r, k; \lambda\}$ supplementary difference sets satisfying the following conditions:

$$v \text{ is an odd positive integer,} \quad (4)$$

$$4(r + k - \lambda) \geq 2, \quad (5)$$

$$2(v - 2(r + k - \lambda)) \geq 1. \quad (6)$$

These supplementary difference sets give matrices M satisfying (1)–(3) (Proposition 6).

This paper is organized as follows. In Section 2, we give definitions and we recall notions on supplementary difference sets and D -optimal designs. Some basic facts on these subjects are also provided. In Section 3, we give some observations on skew-symmetric supplementary difference sets. In Section 4, we describe how to classify skew-symmetric supplementary difference sets satisfying (4)–(6). In Section 5, we give a classification of skew-symmetric 2- $\{v; r, k; \lambda\}$ supplementary difference sets satisfying (4)–(6) for $v \leq 51$ (Theorem 9). This is the main result of this paper. Skew-symmetric circulant D -optimal designs satisfying (9) are corresponding to a special class of supplementary difference sets. In Section 6, as a consequence of Theorem 9, we give a classification of skew-symmetric circulant D -optimal designs meeting (8) for orders up to 110.

2 Preliminaries

In this section, we give definitions and we recall notions on supplementary difference sets and D -optimal designs. Some basic facts on these subjects are also provided.

2.1 Supplementary difference sets

Let $\mathbb{Z}_v = \{0, 1, \dots, v-1\}$ be the ring of integers modulo v , where $v > 2$. For $A \subset \mathbb{Z}_v$ and $i \in \mathbb{Z}_v$, define

$$P_A(i) = |\{(x, y) \in A \times A \mid y - x = i\}| \text{ and} \\ P_A = (P_A(1), P_A(2), \dots, P_A(v-1)).$$

Let A and B be an r -subset and a k -subset of $\mathbb{Z}_v$, respectively. If a pair (A, B) satisfies

$$P_A + P_B = (\lambda, \lambda, \dots, \lambda),$$

then it is called a 2- $\{v; r, k; \lambda\}$ *supplementary difference set*. We refer to [3, 9, 16, 17] for basic facts on supplementary difference sets.

Lemma 2 (Wallis [16, Lemma 1]). *If there exists a 2- $\{v; r, k; \lambda\}$ supplementary difference set, then*

$$r(r-1) + k(k-1) = \lambda(v-1). \quad (7)$$

Chadjipantelis and Kounias [3, Appendix] gave a correspondence between $2\text{-}\{v; r, k; \lambda\}$ supplementary difference sets and pairs of circulant matrices. Let A and B be an r -subset and a k -subset of $\mathbb{Z}_v$, respectively. Let R_1 and R_2 be the circulant $v \times v$ $(1, -1)$ -matrices with first rows $r_1 = (r_{1,1}, r_{1,2}, \dots, r_{1,v})$ and $r_2 = (r_{2,1}, r_{2,2}, \dots, r_{2,v})$, respectively. The correspondence was defined as follows: $r_{1,i+1} = -1$ if $i \in A$, $r_{1,i+1} = 1$ if $i \notin A$ and $r_{2,i+1} = -1$ if $i \in B$, $r_{2,i+1} = 1$ if $i \notin B$.

Lemma 3 (Chadjipantelis and Kounias [3, Appendix]). *A pair (A, B) is a $2\text{-}\{v; r, k; \lambda\}$ supplementary difference set if and only if $R_1 R_1^T + R_2 R_2^T = 4(r + k - \lambda)I + 2(v - 2(r + k - \lambda))J$.*

2.2 D -optimal designs and supplementary difference sets

A D -optimal design of order n is an $n \times n$ $(1, -1)$ -matrix having maximum determinant. Ehlich [4] showed that for $n \equiv 2 \pmod{4}$ and $n > 2$, any $n \times n$ $(1, -1)$ -matrix M satisfies

$$\det M \leq (2n - 2)(n - 2)^{(n-2)/2}, \quad (8)$$

and that equality is possible only if $2n - 2$ is a sum of two perfect squares. Moreover, if $n = 2v \equiv 2 \pmod{4}$, and both R_1 and R_2 are $v \times v$ commutative $(1, -1)$ -matrices such that

$$R_1 R_1^T + R_2 R_2^T = (2v - 2)I + 2J, \quad (9)$$

then

$$X(R_1, R_2) = \begin{pmatrix} R_1 & R_2 \\ -R_2^T & R_1^T \end{pmatrix} \quad (10)$$

is a D -optimal design meeting the above bound (8) [4].

If the matrices R_1 and R_2 in (10) are circulant, then $X(R_1, R_2)$ is called a *circulant D -optimal design* meeting (8) [10]. Most of the known D -optimal designs meeting (8) are circulant (see e.g., [1, 3, 5, 9, 10]). If $X(R_1, R_2)$ is a circulant D -optimal design meeting (8), then it was shown in [3] that

$$(v - 2r)^2 + (v - 2k)^2 = 2n - 2, \quad (11)$$

where r and k are the numbers of -1 's in the first rows of R_1 and R_2 , respectively.

By Lemma 3, we have the following:

Lemma 4 (Chadjipantelis and Kounias [3, Appendix]). *Let A and B be an r -subset and a k -subset of $\mathbb{Z}_v$, respectively. Let R_1 and R_2 be the corresponding circulant $v \times v$ $(1, -1)$ -matrices described in Section 2.1. A pair (A, B) is a 2 - $\{v; r, k; r+k-(v-1)/2\}$ supplementary difference set if and only if $X(R_1, R_2)$ in (10) is a circulant D -optimal design of order $2v$ meeting (8), where r and k are the numbers of -1 's in the first rows of R_1 and R_2 , respectively.*

3 Skew-symmetric supplementary difference sets

Let (A, B) be a supplementary difference set. Let R_1 and R_2 denote the corresponding circulant $v \times v$ $(1, -1)$ -matrices described in Section 2.1. Then we consider the following matrix:

$$X(R_1, R_2) = \begin{pmatrix} R_1 & R_2 \\ -R_2^T & R_1^T \end{pmatrix}. \quad (12)$$

We call (A, B) *skew-symmetric* if the corresponding matrix $X(R_1, R_2)$ in (12) is skew-symmetric. Equivalently, (A, B) is skew-symmetric if A satisfies the condition that $0 \notin A$ and if $i \in A$ then $-i \notin A$. In [2, 15, 18], skew-symmetric 2 - $\{v; (v-1)/2, (v-1)/2; (v-3)/2\}$ supplementary difference sets are called complementary difference sets and these difference sets were used to construct skew-Hadamard matrices.

Lemma 5. *The matrix $X(R_1, R_2)$ in (12) is skew-symmetric if and only if $r_{1,1} = 1$ and $r_{1,i} = -r_{1,v+2-i}$ ($i = 2, 3, \dots, v$). If $X(R_1, R_2)$ is skew-symmetric, then $r = \frac{v-1}{2}$.*

Proof. The elementary proof is omitted. □

By Lemma 3, we have the following:

Proposition 6. *If there exists a skew-symmetric 2 - $\{v; r, k; \lambda\}$ supplementary difference set satisfying (4)–(6). Then there exists a matrix M satisfying (1)–(3) for $(\alpha, \beta) = (4(r+k-\lambda), 2(v-2(r+k-\lambda)))$.*

Now we give a remark on the condition (5) for skew-symmetric 2 - $\{v; r, k; \lambda\}$ supplementary difference sets.

Proposition 7. *Suppose that $k \leq \frac{v-1}{2}$. If there exists a skew-symmetric 2 - $\{v; r, k; \lambda\}$ supplementary difference set (A, B) , then $r+k-\lambda \geq 1$, that is, (A, B) satisfies (5).*

Proof. By Lemma 5, $r = \frac{v-1}{2}$. Hence, it follows from (7) that $r + k - \lambda = \frac{(v+2k)(v-2k)-1+4kv}{4(v-1)}$. From the assumption, $r+k-\lambda > 0$. The result follows. $\square$

For the case $k \in \{0, 1\}$, 2- $\{v; r, k; \lambda\}$ supplementary difference sets are characterized as follows. Although the following characterization is somewhat trivial, it was not explicitly stated in the literature. We give a proof for the sake of completeness.

Proposition 8. *The following statements are equivalent.*

- (1) *There exists a skew-symmetric 2- $\{4m-1; 2m-1, k; m-1\}$ supplementary difference set with $k = 0$ and 1.*
- (2) *There exists a circulant Hadamard 2- $(4m-1, 2m-1, m-1)$ design with incidence matrix M satisfying that $M + M^T + I = J$.*

Proof. Suppose that there exists a skew-symmetric 2- $\{4m-1; 2m-1, k; m-1\}$ supplementary difference set (A, B) with $k \in \{0, 1\}$. Then A is a $(4m-1, 2m-1, m-1)$ -difference set. Let M be an incidence matrix of A . Then M is an incidence matrix of a circulant Hadamard 2- $(4m-1, 2m-1, m-1)$ design. Since A satisfies the condition that if $i \in A$ then $-i \notin A$, M satisfies the condition that $M + M^T + I = J$.

Suppose that there exists a circulant Hadamard 2- $(4m-1, 2m-1, m-1)$ design with incidence matrix M satisfying that $M + M^T + I = J$. By reversing the above argument, a $(4m-1, 2m-1, m-1)$ -difference set A satisfying the condition that if $i \in A$ then $-i \notin A$ is constructed. Then $(A, \emptyset)$ and $(A, \{0\})$ are skew-symmetric 2- $\{v; r, k; \lambda\}$ supplementary difference sets with parameters $(v, r, k, \lambda) = (4m-1, 2m-1, 0, m-1)$ and $(4m-1, 2m-1, 1, m-1)$, respectively. $\square$

Suppose that p is a prime with $p \equiv 3 \pmod{4}$. Then it is well known that there exists a circulant Hadamard 2- $(p, \frac{p-1}{2}, \frac{p-3}{4})$ design with incidence matrix A satisfying that $A + A^T + I = J$ (see e.g., [7, Lemma 7.10]). This implies the existence of skew-symmetric supplementary difference sets with parameters 2- $\{p; \frac{p-1}{2}, 0; \frac{p-3}{4}\}$ and 2- $\{p; \frac{p-1}{2}, 1; \frac{p-3}{4}\}$.

4 Classification method

In this section, we describe how to classify skew-symmetric supplementary difference sets satisfying (4)–(6).

4.1 Equivalent supplementary difference sets

If (A, B) is a supplementary difference set, then the following pairs

$$(E0) \quad (\mathbb{Z}_v \setminus A, B) \text{ and } (A, \mathbb{Z}_v \setminus B),$$

$$(E1) \quad (B, A),$$

$$(E2) \quad (\pm A + a, \pm B + b) \text{ for any } a, b \in \mathbb{Z}_v,$$

$$(E3) \quad (dA, dB) \text{ for any } d \in U(\mathbb{Z}_v)$$

are also supplementary difference sets, where $U(\mathbb{Z}_v) = \{d \in \{1, 2, \dots, v-1\} \mid \gcd(d, v) = 1\}$ and d is regarded as an integer for $\gcd(d, v) = 1$. These supplementary difference sets are called *equivalent* [10].

4.2 Classification method

Let (A, B) be a skew-symmetric 2 - $\{v; r, k; \lambda\}$ supplementary difference set satisfying (4), (6). By Lemma 5, $r = \frac{v-1}{2}$. By (E0), we may assume without loss of generality that $k \leq \frac{v-1}{2}$. We note that (A, B) satisfies (5) by Proposition 7 under this assumption. In addition, if $r = k$, then it follows from (7) that $2(v - 2(r + k - \lambda)) = -2$. Hence, we may assume without loss of generality that

$$k < \frac{v-1}{2} = r. \quad (13)$$

Since A corresponds to a skew-symmetric matrix, there exists $A' \subset \mathbb{Z}'_v$ such that $A = A' \cup \{v - j \mid j \in \mathbb{Z}'_v \setminus A'\}$, where $\mathbb{Z}'_v = \{1, 2, \dots, (v-1)/2\} \subset \mathbb{Z}_v$. By (E2), $(A, B + b)$ is a skew-symmetric supplementary difference set for any $b \in \mathbb{Z}_v$. We classify skew-symmetric 2 - $\{v; (v-1)/2, k; \lambda\}$ supplementary difference sets satisfying (4), (6) by the following steps.

- (i) We calculate $\overline{\mathcal{A}} = \{A' \cup \{v - j \mid j \in \mathbb{Z}'_v \setminus A'\} \mid A' \subset \mathbb{Z}'_v\}$. Then we find $\mathcal{A} = \{A \in \overline{\mathcal{A}} \mid P_A(i) \leq \lambda \text{ for all } i\}$.
- (ii) We calculate $\overline{\mathcal{B}} = \{B \subset \mathbb{Z}_v \mid |B| = k, B \preceq B + b \text{ for any } b \in \mathbb{Z}_v\}$, where $\preceq$ is a natural lexicographic order on k -subsets of $\mathbb{Z}_v$. Then we find $\mathcal{B} = \{B \in \overline{\mathcal{B}} \mid P_B(i) \leq \lambda \text{ for all } i\}$.
- (iii) We construct $\mathcal{AB} = \{(A, B) \in \mathcal{A} \times \mathcal{B} \mid P_A + P_B = (\lambda, \lambda, \dots, \lambda)\}$.

(iv) We classify $\mathcal{AB}$.

In Step (i) (resp. (ii)), we found all $((v-1)/2)$ -subsets (resp. k -subsets) of $\mathbb{Z}_v$ by a computer program implemented in C language using functions from the GNU Scientific Library (GSL) software library, then we output A and $(\lambda, \lambda, \dots, \lambda) - P_A$ (resp. B and P_B) to a file. We sorted the above data by $(\lambda, \lambda, \dots, \lambda) - P_A$ (resp. P_B). We found a pair (A, B) with $(\lambda, \lambda, \dots, \lambda) - P_A = P_B$ in Step (iii). Two skew-symmetric 2 - $\{v; (v-1)/2, k; \lambda\}$ supplementary difference sets (A, B) and (A', B') are equivalent if and only if (A', B') is an element of $\{(\pm dA + a, \pm dB + b) \mid d \in U(\mathbb{Z}_v), a, b \in \mathbb{Z}_v\}$. In Step (iv), for (A, B) and (A', B') , we determined whether there exist $d \in U(\mathbb{Z}_v)$ and $a, b \in \mathbb{Z}_v$ such that $(A', B') = (dA + a, dB + b), (dA + a, -dB + b), (-dA + a, dB + b)$ or $(-dA + a, -dB + b)$. This was done by using the program implemented in C language.

5 Classification of skew-symmetric supplementary difference sets

In this section, we give a classification of skew-symmetric 2 - $\{v; r, k; \lambda\}$ supplementary differences sets satisfying (4)–(6) for $v \leq 51$. This is the main result of this paper. As described in Proposition 6, a skew-symmetric 2 - $\{v; r, k; \lambda\}$ supplementary difference set satisfying (4)–(6), gives a matrix M satisfying (1)–(3) for $(\alpha, \beta) = (4(r + k - \lambda), 2(v - 2(r + k - \lambda)))$.

We call (v, r, k, λ) *feasible parameters* for supplementary difference sets if (v, r, k, λ) satisfies (4)–(6), (7) and (13) (see Proposition 7 for (5)). In Table 1, we list the feasible parameters (v, r, k, λ) for $v \leq 75$.

By an approach given in Section 4, our exhaustive computer search completed a classification of skew-symmetric 2 - $\{v; r, k; \lambda\}$ supplementary difference sets satisfying (4)–(6) for the feasible parameters in Table 1 with $v \leq 51$. We used a computer with CPU Intel(R) Core(TM) i7 4790k, 4 Core.

Theorem 9. *Suppose that $v \leq 51$. If there exists a skew-symmetric 2 - $\{v; r, k; \lambda\}$ supplementary difference sets satisfying (4)–(6), then it is equivalent to one of the supplementary difference sets (A, B) with $v \leq 51$ in Table 3.*

For $v \geq 53$, due to the computational complexity, our exhaustive computer search completed a classification of skew-symmetric 2 - $\{v; r, k; \lambda\}$ sup-

Table 1: Parameters of skew-symmetric supplementary difference sets

(v, r, k, λ)	$N(v, r, k, \lambda)$	(v, r, k, λ)	$N(v, r, k, \lambda)$
(3, 1, 0, 0)	1	(43, 21, 15, 15)	0
(7, 3, 0, 1)	1	(45, 22, 11, 13)	0
(7, 3, 1, 1)	1	(47, 23, 0, 11)	1
(11, 5, 0, 2)	1	(47, 23, 1, 11)	1
(11, 5, 1, 2)	1	(49, 24, 9, 13)	0
(13, 6, 3, 3)	1	(51, 25, 0, 12)	0
(15, 7, 0, 3)	0	(51, 25, 1, 12)	0
(15, 7, 1, 3)	0	(53, 26, 14, 16)	?
(19, 9, 0, 4)	1	(55, 27, 0, 13)	0
(19, 9, 1, 4)	1	(55, 27, 1, 13)	0
(21, 10, 6, 6)	1	(57, 28, 21, 21)	?
(23, 11, 0, 5)	1	(59, 29, 0, 14)	1
(23, 11, 1, 5)	1	(59, 29, 1, 14)	1
(25, 12, 4, 6)	0	(61, 30, 6, 15)	0
(27, 13, 0, 6)	0	(61, 30, 10, 16)	0
(27, 13, 1, 6)	0	(61, 30, 15, 18)	?
(29, 14, 7, 8)	1	(63, 31, 0, 15)	0
(31, 15, 0, 7)	1	(63, 31, 1, 15)	0
(31, 15, 1, 7)	1	(67, 33, 0, 16)	1
(31, 15, 6, 8)	1	(67, 33, 1, 16)	1
(31, 15, 10, 10)	1	(67, 33, 12, 18)	?
(35, 17, 0, 8)	0	(67, 33, 22, 23)	?
(35, 17, 1, 8)	0	(69, 34, 18, 21)	?
(37, 18, 10, 11)	0	(71, 35, 0, 17)	1
(39, 19, 0, 9)	0	(71, 35, 1, 17)	1
(39, 19, 1, 9)	0	(71, 35, 15, 20)	?
(41, 20, 5, 10)	0	(71, 35, 21, 23)	?
(43, 21, 0, 10)	1	(73, 36, 28, 28)	?
(43, 21, 1, 10)	1	(75, 37, 0, 18)	0
(43, 21, 7, 11)	0	(75, 37, 1, 18)	0

plementary difference sets satisfying (4)–(6) for the following feasible parameters:

$$\begin{aligned}
(v, r, k, \lambda) = & (55, 27, 0, 13), (55, 27, 1, 13), (59, 29, 0, 14), \\
& (59, 29, 1, 14), (61, 30, 6, 15), (61, 30, 10, 16), \\
& (63, 31, 0, 15), (63, 31, 1, 15), (67, 33, 0, 16), \\
& (67, 33, 1, 16), (71, 35, 0, 17), (71, 35, 1, 17), \\
& (75, 37, 0, 18), (75, 37, 1, 18).
\end{aligned} \tag{14}$$

The skew-symmetric 2- $\{v; r, k; \lambda\}$ supplementary difference sets (A, B) with parameters (14) are listed in Table 3. For the feasible parameters (v, r, k, λ) given in Table 1, the numbers $N(v, r, k, \lambda)$ of the inequivalent skew-symmetric 2- $\{v; r, k; \lambda\}$ supplementary difference sets are also listed in the table.

6 Classification of skew-symmetric circulant D -optimal designs meeting (8)

Skew-symmetric circulant D -optimal designs $X(R_1, R_2)$ in (10) are corresponding to a certain class of skew-symmetric supplementary difference sets satisfying (4)–(6). According to [10], we say that circulant D -optimal designs meeting (8) are *equivalent* if the supplementary difference sets constructed by Lemma 4 are equivalent. In this section, as a consequence of the previous section, we give a classification of skew-symmetric circulant D -optimal designs meeting (8) for orders up to 110.

Let D be a circulant D -optimal design $X(R_1, R_2)$ in (10) of order $n = 2v$ meeting (8). Here we suppose that r and k are the numbers of -1 's in the first rows of R_1 and R_2 , respectively. If D is skew-symmetric, then $r = \frac{v-1}{2}$ by Lemma 5.

We call (n, r, k) *feasible parameters* for skew-symmetric circulant D -optimal designs if (n, r, k) satisfies $r = \frac{v-1}{2}$ and (11). In Table 2, we list the feasible parameters (n, r, k) for $n \leq 200$.

Let S_3, S_7, S_{13}, S_{21} and S_{31} be the skew-symmetric 2- $\{v; r, k; \lambda\}$ supplementary difference sets in Table 3 with $(v, r, k, \lambda) = (3, 1, 0, 0), (7, 3, 1, 1), (13, 6, 3, 3), (21, 10, 6, 6)$ and $(31, 15, 10, 10)$, respectively. Let $D_6, D_{14}, D_{26}, D_{42}$ and D_{62} be the skew-symmetric circulant D -optimal designs $X(R_1, R_2)$ in (10) of orders 6, 14, 26, 42 and 62 meeting (8), constructed by Lemma 4 from S_3, S_7, S_{13}, S_{21} and S_{31} , respectively. From the classification in Theorem 9, we have the following:

Table 2: Parameters of skew-symmetric circulant D -optimal designs

(n, r, k)	$N(n, r, k)$
$(6, 1, 0)$	1
$(14, 3, 1)$	1
$(26, 6, 3)$	1
$(42, 10, 6)$	1
$(62, 15, 10)$	1
$(86, 21, 15)$	0
$(114, 28, 21)$	?
$(146, 36, 28)$	?
$(182, 45, 36)$	?

Corollary 10. *Suppose that $n \leq 110$. If there exists a skew-symmetric circulant D -optimal design $X(R_1, R_2)$ in (10) of order n meeting (8), then it is equivalent to one of D_6 , D_{14} , D_{26} , D_{42} and D_{62} .*

The numbers $N(n, r, k)$ of the inequivalent skew-symmetric circulant D -optimal designs meeting (8) are also listed in Table 2 for the feasible parameters (n, r, k) .

A classification of circulant D -optimal designs $X(R_1, R_2)$ in (10) meeting (8) was given in [10] for orders $n \leq 58$ and $n = 66$, and in [1] for orders $n = 62, 74$ (see [1] for the revised classification for order 26). Our computer search found that D_n ($n = 6, 14, 26$) is equivalent to the circulant D -optimal design, which is constructed by Lemma 4 from the first supplementary difference set given in [10, Table 1], D_{42} is equivalent to the circulant D -optimal design, which is constructed by Lemma 4 from the 19th supplementary difference set given in [10, Table 1], and D_{62} is equivalent to the circulant D -optimal design, which is constructed by Lemma 4 from the 50th supplementary difference set given in [1, Appendix].

Acknowledgment. The authors would like to thank the anonymous referees for helpful comments. This work is supported by JSPS KAKENHI Grant Numbers 15K04976, 26610032.

References

- [1] M. Araya and M. Harada, Classification of circulant D -optimal designs of orders 62 and 74, *J. Statist. Appl.* **4** (2009), 301–312.
- [2] D. Blatt and G. Szekeres, A skew Hadamard matrix of order 52, *Canad. J. Math.* **21** (1969), 1319–1322.
- [3] Th. Chadjiapantelis and S. Kounias, Supplementary difference sets and D -optimal designs for $n \equiv 2 \pmod{4}$, *Discrete Math.* **57** (1985), 211–216.
- [4] H. Ehlich, Determinantenabschätzungen für binäre Matrizen, *Math. Z.* **83** (1964), 123–132.
- [5] R.J. Fletcher, C. Koukouvinos and J. Seberry, New skew-Hadamard matrices of order 4·59 and new D -optimal designs of order 2·59, *Discrete Math.* **286** (2004), 251–253.
- [6] A.V. Geramita and J. Seberry, *Orthogonal Designs, Quadratic Forms and Hadamard Matrices*, Marcel Dekker Inc., New York, 1979.
- [7] A.S. Hedayat, N.J.A. Sloane and J. Stufken, *Orthogonal Arrays*, Springer-Verlag, New York, 1999.
- [8] S.J. Kirkland and B.L. Shader, Tournament matrices with extremal spectral properties, *Linear Algebra Appl.* **196** (1994), 1–17.
- [9] C. Koukouvinos, S. Kounias and J. Seberry, Supplementary difference sets and optimal designs, *Discrete Math.* **88** (1991), 49–58.
- [10] S. Kounias, C. Koukouvinos, N. Nikolaou and A. Kakos, The nonequivalent circulant D -optimal designs for $n \equiv 2 \pmod{4}$, $n \leq 54$, $n = 66$, *J. Combin. Theory Ser. A* **65** (1994), 26–38.
- [11] H. Nozaki and S. Suda, A characterization of skew Hadamard matrices and doubly regular tournaments, *Linear Algebra Appl.* **437** (2012), 1050–1056.
- [12] H. Nozaki and S. Suda, Complex spherical codes with two inner products, *European J. Combin.* **51** (2016), 511–518.

- [13] K.B. Reid and E. Brown, Doubly regular tournaments are equivalent to skew Hadamard matrices, *J. Combin. Theory Ser. A* **12** (1972), 332–338.
- [14] P. Rowlinson, On 4-cycles and 5-cycles in regular tournaments, *Bull. London Math. Soc.* **18** (1986), 135–139.
- [15] G. Szekeres, Tournaments and Hadamard matrices, *Enseignement Math. (2)* **15** (1969), 269–278.
- [16] J. (Seberry) Wallis, On supplementary difference sets, *Aequationes Math.* **8** (1972), 242–257.
- [17] J. (Seberry) Wallis, A note on supplementary difference sets, *Aequationes Math.* **10** (1974), 46–49.
- [18] A.L. Whiteman, An infinite family of skew Hadamard matrices, *Pacific J. Math.* **38** (1971), 817–822.

Table 3: Skew-symmetric supplementary difference sets (A, B)

14

$(v, r, k, \lambda) = (3, 1, 0, 0)$ $A = \{2\} \ B = \emptyset$
$(v, r, k, \lambda) = (7, 3, 0, 1), (7, 3, 1, 1)$ $A = \{3, 5, 6\} \ B = \emptyset, \{0\}$
$(v, r, k, \lambda) = (11, 5, 0, 2), (11, 5, 1, 2)$ $A = \{2, 6, 7, 8, 10\} \ B = \emptyset, \{0\}$
$(v, r, k, \lambda) = (13, 6, 3, 3)$ $A = \{4, 7, 8, 10, 11, 12\} \ B = \{0, 2, 8\}$
$(v, r, k, \lambda) = (19, 9, 0, 4), (19, 9, 1, 4)$ $A = \{2, 3, 8, 10, 12, 13, 14, 15, 18\} \ B = \emptyset, \{0\}$
$(v, r, k, \lambda) = (21, 10, 6, 6)$ $A = \{2, 3, 9, 11, 13, 14, 15, 16, 17, 20\} \ B = \{0, 1, 7, 9, 12, 17\}$
$(v, r, k, \lambda) = (23, 11, 0, 5), (23, 11, 1, 5)$ $A = \{5, 7, 10, 11, 14, 15, 17, 19, 20, 21, 22\} \ B = \emptyset, \{0\}$
$(v, r, k, \lambda) = (29, 14, 7, 8)$ $A = \{4, 5, 6, 8, 9, 10, 12, 13, 15, 18, 22, 26, 27, 28\} \ B = \{0, 1, 11, 13, 15, 18, 21\}$
$(v, r, k, \lambda) = (31, 15, 0, 7), (31, 15, 1, 7)$ $A = \{3, 6, 11, 12, 13, 15, 17, 21, 22, 23, 24, 26, 27, 29, 30\} \ B = \emptyset, \{0\}$
$(v, r, k, \lambda) = (31, 15, 6, 8)$ $A = \{3, 6, 11, 12, 13, 15, 17, 21, 22, 23, 24, 26, 27, 29, 30\} \ B = \{0, 1, 15, 20, 22, 28\}$
$(v, r, k, \lambda) = (31, 15, 10, 10)$ $A = \{4, 6, 7, 12, 16, 17, 18, 20, 21, 22, 23, 26, 28, 29, 30\} \ B = \{0, 1, 4, 5, 8, 11, 16, 18, 20, 29\}$
$(v, r, k, \lambda) = (43, 21, 0, 10), (43, 21, 1, 10)$ $A = \{2, 3, 5, 7, 8, 12, 18, 19, 20, 22, 26, 27, 28, 29, 30, 32, 33, 34, 37, 39, 42\} \ B = \emptyset, \{0\}$
$(v, r, k, \lambda) = (47, 23, 0, 11), (47, 23, 1, 11)$ $A = \{5, 10, 11, 13, 15, 19, 20, 22, 23, 26, 29, 30, 31, 33, 35, 38, 39, 40, 41, 43, 44, 45, 46\} \ B = \emptyset, \{0\}$
$(v, r, k, \lambda) = (59, 29, 0, 14), (59, 29, 1, 14)$ $A = \{2, 6, 8, 10, 11, 13, 14, 18, 23, 24, 30, 31, 32, 33, 34, 37, 38, 39, 40, 42, 43, 44, 47, 50, 52, 54, 55, 56, 58\} \ B = \emptyset, \{0\}$
$(v, r, k, \lambda) = (67, 33, 0, 16), (67, 33, 1, 16)$ $A = \{2, 3, 5, 7, 8, 11, 12, 13, 18, 20, 27, 28, 30, 31, 32, 34, 38, 41, 42, 43, 44, 45, 46, 48, 50, 51, 52, 53, 57, 58, 61, 63, 66\} \ B = \emptyset, \{0\}$
$(v, r, k, \lambda) = (71, 35, 0, 17), (71, 35, 1, 17)$ $A = \{7, 11, 13, 14, 17, 21, 22, 23, 26, 28, 31, 33, 34, 35, 39, 41, 42, 44, 46, 47, 51, 52, 53, 55, 56, 59, 61, 62, 63, 65, 66, 67, 68, 69, 70\} \ B = \emptyset, \{0\}$